

(1830); Hook. Fl. ii. 82: *Eritrichium oxycaryum*, Gray, Proc. Am. Acad. x. 58 (1874), and Syn. Fl. 193: *Krynitzkia oxycarya*, Gray, l. c.

25. *C. MICROSTACHYS* = *Krynitzkia microstachys*, Greene in Gray, Proc. Am. Acad. xx. 269, and Syn. Fl. Suppl. 425.

26. *C. ROSTELLATA* = *Krynitzkia rostellata*, Greene, Bull. Cal. Acad. i. 203; Gray, Syn. Fl. Suppl., l. c.

27. *C. SPARSIFLORA* = *Krynitzkia sparsiflora*, Greene, l. c., and Gray l. c.

28. *C. RAMOSISSIMA* = *Krynitzkia ramosissima*, Greene, Bull. Cal. Acad. i. 203; Gray, Suppl. 428 and, in part, of Proc. Am. Acad. xx. 277.

29. *C. GLOMERIFLORA*. Annual, 2—4 inches high, diffusely branching and flowering from the base, very hispid throughout; leaves linear-oblong, $\frac{1}{4}$ — $\frac{1}{2}$ inch long: flowers in glomerules of 2 or 3 in the axils of the leaves and at the ends of the branchlets: corolla very minute: calyx very bristly, its linear segments only $\frac{1}{2}$ line long, a little surpassed by the ovate-acuminate speckled nutlet, whose ventral groove is closed