

30. *C. CEDROSENSIS* = *Krynitzkia Cedrosensis*, Greene, l. c. 204.

31. *C. MARITIMA* = *K. maritima*, Greene, l. c.

32. *C. CLEVELANDI*. A foot or more in height, with few ascending branches rather rigid, and bearing two or three short racemes at summit; hispid throughout with slender but rigid pungent bristles, and without appressed pubescence: calyx slender, appressed to the rachis (as in *C. flaccida*): nutlets 2 or 1.

Common in shaded places along streamlets in the hills back of San Diego, where it was collected by Mr. Cleveland and the writer in April, 1885, the specimens having been largely distributed by me as "*K. microstachys*, Greene," from which it is very distinct, being as it were intermediate between that and *C. leiocarpa*, but with more slender nutlets than those of that species. It was also obtained, in the same year at All Saint's Bay, Lower California

++ ++ Nutlets four.

33. *C. LEIOCARPA* = *Echinosperrum leiocarpum*, Fisch. & Mey. Sem. Hort. Petrop. 1835. 36, also in Linnæa (Lit. Bericht), 1837. 104; Don. Gen. Syst. iv. 373: *Krynitzkia leiocarpa*, Fisch. & Mey. op. cit. 1841. 52: A. DC. Prodr. x. 134; Gray, l. c.—Six inches to a foot high, diffusely branched, canescent with an appressed pubescence, and with more or less of pilose-hispid spreading hairs: inflorescence short-spicate or somewhat glomerate and leafy: calyx a line long, the segments not much elongated or attenuate above the nutlets, the latter with closed groove which is not forked at base.

Common in the sand hills of San Francisco, well out on California street, April, 1886; also obtained at Point Reyes by Mrs. Curran, and near Gilroy by Mr. Hickman. Plant seldom much hispid except upon the calyx, and, in the San Francisco