

**K. stipitata**, Greene. Ten to 18 inches high, erect and simple or with ascending branches from the base, light green, nearly glabrous: calyx nearly sessile, segments spreading foliaceous, in fruit often 6 lines long: corolla short-funnelform, 3-6 lines broad: nutlets slender-ovoid, the back covered with blunt tubercles and transverse wrinkles, divergent, stipitate. This, according to Greene, is one of the most common species in Central California. Moist land.

**K. Hickmanii**, Greene. Very slender, diffuse, the filiform racemose branches 6-10 inches long: calyx a line long, on filiform pedicels: corolla a line or more broad: nutlets dark colored, tuberculate. Monterey Co.

**K. hirta**, Greene. Annual, more than a foot high, erect, flaccid, simple below with many pairs of connate-sheathing linear leaves, loosely racemose above, bristly hairy: racemes in pairs: pedicels slender, a line long: calyx lobes erect, very hairy: corolla 3 lines broad: ovoid nutlets dark colored, the back granulate and obscurely wrinkled. Umpqua Valley, Or.

**K. scripta**, Greene. Somewhat succulent strigose-hairy: branches prostrate, 6-10 inches long: pedicels stout in the axils of leafy bracts: sepals oblanceolate at length, standing vertically in row: nutlets a line long, the back dark and smooth, marked by a sharp irregular flexuose with white ridges, these beset with tufts of short spreading bristles.

Section 3 is equivalent to **Cryptanthe** in *Bay-Reg. Bot.* The following species are to be added:

**C. flaccida**, Greene, is **K. oxycarya** of this book.

**K. Clevelandi**, Greene. A foot or more high with few ascending branches rough with bristly hairs: calyx slender, appressed to the rachis: nutlets 2 or 1, shining.

**K. hispidissima**, Greene. Size of the last, but more branching, more densely hispid with softer hairs, and the inflorescence more elongated: corolla conspicuous: nutlets 4, similar to those of *K. leiocarya*, much surpassed by the slender calyx lobes. San Luis Obispo Co.

**K. nemaclada**, Greene. Slender, very diffusely branching, a foot high, sparsely bristly-hairy: spikes very loose, almost filiform: calyx a line long appressed to the rachis, bristly: nutlets 4, ovoid-acuminate  $\frac{1}{2}$  a line long, shining. Colusa Co.

**K. Rattani**, Greene. Hispid with slender hairs and slightly canescent, about a foot high, slender but rigid: leaves linear: spikes in 3s on an elongated naked common peduncle, rather densely flowered: calyx appressed to the rachis, its bristles spreading and straight: nutlets (3 maturing) lance-ovoid  $1\frac{1}{2}$  lines long, brownish and smooth but not shining. First collected by Hickman in Monterey Co.

**K. crinita**, Greene. Annual, slender 8-12 inches high, somewhat fastigiately branched from the base, rather stiffly hirsute: leaves linear: dense spikes elongated: